

# The Association of Multimorbidity With Whole Health Activities Among Adults in the United States: Evidence From the NHIS and BRFSS

Global Advances in Integrative Medicine and Health

Volume 12: 1–9


© The Author(s) 2023

Article reuse guidelines:

[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)

DOI: 10.1177/27536130231156857

[journals.sagepub.com/home/gam](https://journals.sagepub.com/home/gam)

Rolake A. Neba, PharmD<sup>1</sup> , Mayela Warner, PharmD<sup>1</sup>, Sydney E. Manning, MPH<sup>1</sup>, R. Constance Wiener, MA, DMD, PhD<sup>2</sup>, and Usha Sambamoorthi, PhD<sup>1</sup>

## Abstract

**Background:** Whole health is a holistic approach encompassing integrative medicine, emotional, and spiritual health and is critical to improving health outcomes among individuals with multimorbidity.

**Objective:** To examine the prevalence of Whole Health activities and the association of multimorbidity and Whole Health activities using nationally representative datasets.

**Methods:** As no single dataset has information on Whole Health self-care activities, data from the 2017 National Health Interview Survey (n = 25 134) was used to measure participants' mind-body therapy usage, sleep, mental health, and physical activity. We used the 2017 Behavioral Risk Factor Surveillance System (n = 347 029) to assess regular vegetable and/or fruit consumption.

**Results:** A significantly lower percentage of adults with multimorbidity had adequate sleep (58.2%vs.67.1%), no psychological distress (71.8%vs.82.1%), adequate physical activity (48.2%vs.62.1%), and regular vegetable and/or fruit consumption (54.2% vs.56.6%) compared to those without multimorbidity. Although lower percentages of adults with multimorbidity utilized mind-body therapies (22.9%vs.25.2%), the association was reversed when adjusted for socioeconomic factors. In the fully adjusted models, adults with multimorbidity were more likely to use mind-body therapies (AOR = 1.19, 95%CI = 1.09, 1.31). Furthermore, when adjusting for other independent variables, the associations of multimorbidity with sleep, psychological distress, and diet were exacerbated, and the association of multimorbidity with physical activity was attenuated.

**Conclusion:** Adults with multimorbidity were less likely to engage in most of the Whole Health activities except mind-body therapies compared to the no multimorbidity group. Findings suggest that adjustment for other factors such as age and socioeconomic status changed the magnitude and direction of the association of multimorbidity with Whole Health activities.

## Keywords

whole health, diet, mental health, sleep, physical activity, multimorbidity

Received August 23, 2022; Revised December 20, 2022. Accepted for publication January 24, 2023

<sup>1</sup>Department of Pharmacotherapy, College of Pharmacy, University of North Texas Health Science Center, Fort Worth, TX, USA

<sup>2</sup>Department of Dental Public Health and Professional Practice, School of Dentistry, West Virginia University, Morgantown, WV, USA

## Corresponding Author:

Rolake A. Neba, Department of Pharmacotherapy, College of Pharmacy, University of North Texas Health Science Center, Fort Worth, TX 76107, USA.

Email: [rolake.neba@unthsc.edu](mailto:rolake.neba@unthsc.edu)



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use,

reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and

Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

## Introduction

Multimorbidity, the concurrent presence of 2 or more chronic conditions, is a global concern for prevention and management, as multimorbidity is associated with health decline and mortality.<sup>1</sup> The prevalence of multimorbidity in the U.S. is approximately 1 in 4 U.S. adults.<sup>1</sup> For individuals with multimorbidity, the implementation of disease-specific guidelines to improve health has fallen short. Clinical guidelines and quality standards derived from clinical trials have limited relevance for those with multimorbidity because trials typically exclude individuals with multiple chronic conditions.<sup>2</sup> For individuals with multiple conditions, challenges exist to implementing evidence-based practices due to the proliferation of the standard of care for each disease.<sup>3</sup> Health systems, providers, patients, government agencies, and other stakeholders are working towards more sustainable models of care.

One of the sustainable care models that may improve health among those with multimorbidity is the Whole Health perspective. The Whole Health perspective not only focuses on singular, specific diseases and disease guidelines but also on “what matters to the person.” The Whole Health model is based on the Institute of Medicine and the World Health Organization (WHO) standards, in which balanced health involves health and healing approaches to mental and emotional; physical and structural; nutritional and chemical; environmental, and spiritual aspects of care.<sup>4</sup>

Individuals from various backgrounds and disciplines such as pharmacists, psychologists, social workers, clinicians, chaplains, physical assistants, complementary health practitioners, and many others, have contributed to the evolution of the Whole Health Model.<sup>5</sup> There are different versions of the Whole Health approach from the National Institute of Whole Health (NIWH),<sup>6</sup> the National Center for Complementary and Integrative Health (NCCIH),<sup>4</sup> the Veterans Health Administration (VA),<sup>7</sup> the Whole Health Institute,<sup>8</sup> and TakeCare<sup>9</sup> that extended the NIWH model. However, all these versions have some common elements such as person-centeredness, disease-agnostic holistic approach, integration of traditional and complementary therapies; self-care as a pillar of health and well-being; the role of social determinants of health (SDoH), and care supported by health systems, programs, and policies.<sup>5-9</sup>

It is essential to have an evidence-based approach for Whole Health practices to contain costs, coordinate care, satisfy the patients, and improve outcomes,<sup>10</sup> specifically among those with multimorbidity.<sup>11</sup> There is some evidence that Whole Health activities that intersect self-care activities and integrated care with both conventional and complementary medicine can improve clinical outcomes,<sup>12</sup> decrease costs,<sup>13</sup> well-being of employees,<sup>13</sup> patient satisfaction,<sup>13</sup> physical and emotional health,<sup>12</sup> and employment.<sup>12</sup>

For many health conditions, self-care in terms of Whole Health activities can play a significant role in primary,

secondary, and tertiary prevention activities.<sup>14</sup> All types of prevention activities can succeed if the 5 critical activities of self-care are practiced. These activities are (1) the power of mind, (mind-body therapies), (2) recharge (adequate sleep), (3) mental health (no or low psychological distress), (4) moving the body (adequate physical activity), and (5) nutrition (regular vegetable and/or fruit consumption).<sup>15</sup>

The growing burden of multimorbidity coupled with information technology has propelled individuals towards high-value practices such as Whole Health activities.<sup>16</sup> However, limited literature is available on the prevalence of Whole Health activities, specifically among individuals with multimorbidity who need it the most. We acknowledge that there have been studies on the reciprocal relationships of individual activities such as sleep,<sup>17</sup> physical activity,<sup>18,19</sup> diet,<sup>20</sup> and psychological distress<sup>21</sup> with incident multimorbidity and the association of type of multimorbidity on use of alternative and complementary medicine including mind-body therapies.<sup>22</sup> However, studies on the prevalence rates of all Whole Health activities among those with multimorbidity are sparse. Therefore, this research aimed to examine the prevalence of Whole Health activities and the association of multimorbidity and Whole Health activities among adults in the U.S. using nationally representative datasets.

## Methods

### Study Design and Data Sources

We used a cross-sectional design and analyzed data on adults aged 18 or older from 2 nationally representative health surveys. As no 1 health survey collected information on all 5 self-care activities of the Whole Health model, we used the 2017 National Health Interview Survey (NHIS) to measure the prevalence of mind-body therapies, adequate sleep, psychological stress, and physical activity, and the 2017 Behavioral Risk Factor Surveillance System (BRFSS) to assess regular vegetable and/or fruit consumption. We selected 2017 because it represented the most recent available data on Whole Health self-care activities.

We harmonized measures in the 2 datasets, and variables were recoded to have the same categories or levels between the NHIS and the BRFSS. The study samples were restricted to adults without missing data on Whole Health self-care activities and multimorbidity variables and without missing age (BRFSS 2017). The analytical samples comprised 25, 133 adults from the NHIS and 343 341 adults from the BRFSS, representing about 231.8 million and 183 million adults, respectively.

## Measures

### Dependent Variables

*NHIS-2017 - Mind-body therapies (Yes/No).* Adults who practiced mantra, mindfulness, and spiritual meditation<sup>23</sup> or

meditation through Yoga, Tai chi, Qigong, or guided imagery during the past 12 months were considered to use mind-body therapies.

**NHIS 2017 - Adequate Sleep (Yes/No).** We defined adequate sleep using the duration of sleep based on the consensus statement of the American Academy of Sleep Medicine and Sleep Research Society.<sup>24</sup> Based on the question “On average, how many hours of sleep do you get in a 24-hour period?”, we classified adults with at least 7 hours of sleep per night as having adequate sleep.

**NHIS 2017 - Adequate Physical Activity (Yes/No).** Adequate physical activity was derived from questions about frequency, duration, and intensity of leisure time physical activity and strength training reported weekly. Using the 2008 Physical Activity Guidelines for Americans,<sup>25</sup> adults with at least 150 minutes of moderate-intensity activity per week or who engaged in muscle-strengthening activities at least 2 times per week were considered to have adequate physical activity.<sup>26</sup>

**NHIS 2017 – Low or No Psychological Distress (Yes/No).** A person’s mental health was measured by the Kessler Psychological Distress Scale (K6). The K6 consists of 6 questions on nervousness, hopelessness, worthlessness, restless/fidgety, sadness not able to be alleviated, and struggling in effort during the past 30 days.<sup>27</sup> Each question is scored from 0 (None of the time) to 4 (All of the time) and the sum of these scores range from 0 to 24. Based on a study that validated a sub-threshold cut point for moderate distress, we classified adults with scores less than 5 as having low psychological distress.<sup>28</sup>

**BRFSS 2017 - Regular vegetable and/or fruit consumption (Yes/No).** As of 2017, BRFSS now asks 6 questions as part of their dietary module.<sup>29</sup> These questions ask how frequently a respondent consumed different types of fruits and/or vegetables in the previous 30 days. In the absence of more comprehensive dietary questions, this study defined individuals with “regular vegetable and/or fruit consumption” as those who had at least 1 fruit and vegetable per day in the past 30 days.

### **Key Independent Variable: Multimorbidity (Yes/No)**

This article defined multimorbidity as the co-occurrence of 2 or more chronic conditions in a single individual.<sup>30</sup> Multimorbidity was measured with an indicator variable 1 representing the presence of multimorbidity and 0 representing no multimorbidity. Adults with 2 or more of the following chronic conditions in BRFSS and NHIS (arthritis, asthma, cancer, chronic kidney disease, chronic obstructive pulmonary disease, diabetes, hyperlipidemia, hypertension, heart disease, and stroke) were classified as

having multimorbidity. These chronic conditions were based on the availability of data as well as the framework for managing multiple chronic conditions for research, program, and policy.<sup>31</sup>

### **Other Explanatory Variables**

For adjusted analyses, we controlled for biologic factors, socioeconomic factors, and lifestyle factors. Biological factors included: sex (male/female); age in years (18-39, 40-49, 50-64, 65+); and race/ethnicity (White, African American, Hispanic/Latino, Other). Socioeconomic factors included: education (less than high school, high school, some college, college); health insurance status (yes, no); poverty relative to the federal poverty line (FPL) for NHIS (<100% FPL, 100-200% FPL, 200-400% FPL, >400% FPL); and annual household income for BRFSS (less than \$10k, \$10k-25k, \$25-75k, >\$75k). Social support included marital status (married, never married, divorced/widowed/separated). Lastly, lifestyle factors included obesity (under/normal weight, overweight, obese); smoking status (current, former, never); and alcohol use, which differed between surveys. For NHIS, alcohol use was measured as former, current, or never, whereas for BRFSS, it was measured as heavy drinking, moderate drinking, or no drinking. We also included a variable for the respondent’s region (Northeast, Midwest, South, West). We included a missing indicator for variables with missing data. Although missing data were included in the analysis, we did not present them in tables.

### **Statistical Analyses**

Statistically significant group differences in Whole Health self-care activities by multimorbidity status were tested with Rao-Scott chi-square tests. Adjusted relationships of multimorbidity with Whole Health activities were tested with separate multivariable logistic regressions on each of the Whole Health activities. In these logistic regressions, we controlled for age, sex, race/ethnicity, education, income, health insurance, obesity, smoking, and region. We conducted the regression analysis using multiple models to examine changes in the association of multimorbidity with Whole Health activities. The first model included only multimorbidity (Yes/No) as an independent variable. In the second model, we adjusted for multimorbidity, sex, age, race and ethnicity, education, and poverty status. The third model was fully adjusted for multimorbidity, sex, age, race and ethnicity, education, poverty status, health insurance, marital status, body mass index, current smoking, alcohol use, and region.

To account for the complex survey design of the NHIS, we used SAS survey procedures with clustering, strata, and weights.<sup>32</sup> Statistical analysis with BRFSS used the final weights provided in the dataset to generalize from the sample to the population.<sup>33</sup>

**Table 1.** Description of Selected Characteristics of Respondents. Adults (age ≥ 18 Years). National Health Interview Survey (NHIS) 2017 and Behavioral Risk Factor Surveillance System (BRFSS) 2017.

	NHIS		BRFSS	
	N	Wt.%	N	Wt.%
All	25 133	100	343 341	100
Multimorbidity				
Yes	10 502	36.0	176 655	40.8
No	14 631	64.0	170 374	59.2
Sex				
Female	13 772	51.8	193 911	51.9
Male	11 361	48.2	149 430	48.1
Age groups				
18-39 Years	8 134	38.5	69 155	35.2
40-49 Years	3 634	16.1	42 599	15.6
50-64 Years	6 535	25.6	106 875	27.3
65 years or older	6 830	19.8	124 712	21.9
Race and ethnicity				
White	17 765	64.8	264 915	64.4
African American	2 737	12.0	25 097	10.7
Hispanic/Latino	3 052	16.0	26 012	15.9
Other race	1 579	7.1	22 489	7.5
Marital status				
Married	12 634	60.3	192 729	56.2
Div/Sep/Wid	6 772	16.9	96 782	20.6
Never married	5 687	22.6	52 512	22.8
Education				
Less than HS	2 769	11.6	22 292	12.5
High school	5 999	23.9	90 794	27.6
Some college	7 840	30.3	96 685	31.9
College	8 455	33.9	132 939	27.8
Poverty status				
<100% FPL	3 234	10.9	32 636	12.4
100-<200%	4 238	15.8	57 013	15.7
200-<400%	6 715	26.9	93 729	25.9
≥ 400%	9 520	40.4	79 048	18.5
Health insurance				
Insured	22 823	90.0	317 460	88.4
Not insured	2 233	10.0	24 885	11.2
Used mind-body therapies				
Yes	6 347	24.4	—	—
No	18 786	75.6	—	—
Adequate sleep				
Yes	16 043	63.9	—	—
No	9 090	36.1	—	—
Adequate physical activity				
Yes	14 104	57.1	—	—
No	11 029	42.9	—	—
Low or No psychological distress				
Yes	19 580	78.4	—	—
No	5 553	21.6	—	—
Number of Whole Health activities met				
None	1 210	4.6	—	—
One	4 554	17.7	—	—

(continued)

**Table 1.** (continued)

	NHIS		BRFSS	
	N	Wt.%	N	Wt.%
Two	9 069	36.7	—	—
Three	7 818	31.3	—	—
All 4	2 482	9.7	—	—
Regular vegetable and/or fruit consumption				
Yes	—	—	201 284	55.6
No	—	—	142 057	44.4

Based on 25 133 adult respondents (age 18 years or older) from the National Health Interview Survey and 343 341 adult respondents from the Behavioral Risk Factor Surveillance System. Due to missing data in BRFSS 2017 (Race and Ethnicity, Marital status, Education, and health insurance) and NHIS 2017 (Marital status, Education, and health insurance), the percentages may not total 100%.

Div/Sep/Wid: Divorced/Separated/Widowed; FPL: Federal Poverty Line; N: Unweighted number of individuals in a category. Wt. %: Weighted; HS: High school.

## Results

**Table 1** summarizes respondent characteristics in terms of sex, age, race and ethnicity, education, marital status, and health insurance in the BRFSS and NHIS. In general, the characteristics were similar between the NHIS and the BRFSS. The prevalence rates of Whole Health activities were: 24.4% (mind-body therapies), 63.9% (adequate sleep), 78.4% (no or low psychological distress), 57.1% (adequate physical activity), and 55.6% (regular vegetable and/or fruit consumption).

### Multimorbidity and Whole Health Activities

We observed statistically significant differences ( $P < .001$ ) in all measures of Whole Health activities by multimorbidity status (**Table 2**). A lower percentage of adults with multimorbidity used mind-body therapies (22.9% vs 25.5%), had adequate sleep (58.2% vs 67.1%), had adequate physical activity (48.2% vs 62.1%), low to no psychological distress (71.8% vs 82.1%), and reported having a regular vegetable and/or fruit consumption (54.2% vs 56.6%) compared to those without multimorbidity. Based on NHIS, only 7.1% of those with and 11.1% without multimorbidity reported Whole Health in all 4 Whole Health activities.

In unadjusted logistic regressions (**Table 3**), individuals with multimorbidity were significantly less likely than those without multimorbidity to use mind-body therapies (UOR = .87, 95%CI .81-.94), get adequate sleep (OR = .68, 95%CI .64-.73), have no or low psychological distress (UOR = .56, 95%CI .51-.60), get adequate physical activity (UOR = .56, 95%CI .53-.60), and have regular vegetable and/or fruit consumption (UOR = .91, 95%CI = .88, .94).

Fully adjusted multivariable regression analyses in **Table 3** revealed similar results for adequate sleep (AOR = .64, 95%

**Table 2.** Prevalence of Whole Health Activities by Multimorbidity Status. Among Adults (age ≥ 18 years). National Health Interview Survey (NHIS) 2017 and Behavioral Risk Factor Surveillance System (BRFSS) 2017.

Variable	Multimorbidity		No Multimorbidity		Chi-sq	P-Value
	N	Wt.%	N	Wt.%		
		NHIS 2017				
NHIS ALL	10 502	<b>36.0</b>	<b>14 631</b>	<b>64.0</b>		
Used mind-body therapies					10.336	<.01
Yes	2429	22.9	3918	25.2		
No	8073	77.1	10 713	74.8		
Low or No psychological distress					252.822	<.001
Yes	7606	71.8	11 974	82.1		
No	2896	28.2	2657	17.9		
Adequate sleep					143.245	<.001
Yes	6237	58.2	9806	67.1		
No	4265	41.8	4825	32.9		
Adequate physical activity					281.606	<.001
Yes	4889	48.2	9215	62.1		
No	5613	51.8	5416	37.9		
Number of Whole Health activities met					532.481	<.001
None	809	7.9	401	2.8		
One	2393	22.6	2161	14.9		
Two	3870	37.0	5199	36.5		
Three	2692	25.4	5126	34.7		
All four	738	7.1	1744	11.1		
		BRFSS 2017				
BRFSS ALL	176 655	40.8	170 374	59.2		
Regular vegetable and/or fruit consumption					39.108	<.001
Yes	100 075	54.2	101 209	56.6		
No	74 824	45.8	67 233	43.4		

Based on 25 133 adult respondents (age 18 years or older) from NHIS and 343 341 adult respondents from the BRFSS. Mind-body therapies, low or no psychological distress, adequate sleep, adequate physical activity, and the number of Whole Health self-care activities were measured using NHIS 2017, and regular vegetable and/or fruit consumption was measured using BRFSS 2017. Statistical significance is based on Rao-Scott Chi-square tests.

N: Unweighted number of individuals in a category. Wt. %: Weighted column percent.

CI = .59-.69), no or low psychological distress (AOR = .41, 95% CI = .37-.45), adequate physical activity (AOR = .82, 95% CI .76-.88), and regular vegetable and/or fruit consumption (AOR = .91, 95% CI = .88-.94). However, the adults with multimorbidity had higher odds of mind-body therapy use in the fully adjusted model (AOR = 1.19, 95% CI = 1.09, 1.31). An examination of model 2 suggests that adjustment for education reversed the association, and adults with multimorbidity were more likely to use mind-body therapies (AOR = 1.11, 95% CI 1.01-1.21).

## Discussion

The power of Whole Health activities in terms of self-care as 1 of the pillars of a Whole Health model has been well documented in the scientific literature.<sup>34</sup> In this article, we examined the prevalence and association of multimorbidity with Whole Health activities as part of a Whole Health model. The prevalence rates found in our study were consistent with those from prior published studies that have separately

analyzed self-care activities. For example, Malecki et al, described that physical inactivity in the United States ranges from 17.3% to 47.7%,<sup>35</sup> while the CDC states 31.6% of adults get insufficient sleep according to NHIS data from 2011-2014.<sup>36</sup>

In fully adjusted analyses, adults with multimorbidity were found to have lower odds of Whole Health activities compared to the no multimorbidity group except for mind-body therapies. Our study findings also highlight how the adjustment of other independent variables changed the strength and directions of associations of multimorbidity with the type of Whole Health activities. For example, adjusting for age and socioeconomic status, the association of multimorbidity with sleep, psychological distress, and diet was exacerbated. However, the same factors attenuated the association of multimorbidity with physical activity and changed the direction of the association with mind-body therapies. These findings suggest that socioeconomic factors differentially contribute to the association of multimorbidity with Whole Health activities. Therefore, programs

**Table 3.** Odds Ratios and 95% Confidence Intervals (CI). From Separate Logistic Regressions on Whole Health Activities. Adults (age ≥ 18 years). National Health Interview Survey (NHIS) 2017 and Behavioral Risk Factor Surveillance System (BRFSS) 2017.

Variable	Model 1	Model 2	Model 3
	UOR (95% CI)	AOR (95% CI)	AOR (95% CI)
NHIS 2017			
Dependent variable: Mind-body therapies used			
Multimorbidity			
Yes	.87 (.81-.94)	1.11 (1.01-1.21)	1.19 (1.09-1.31)
No (Reference)			
Dependent variable: Adequate sleep			
Multimorbidity			
Yes	.68 (.64-.73)	.60 (.55-.65)	.64 (.59-.69)
No (Reference)			
Dependent variable: Low or No psychological distress			
Multimorbidity			
Yes	.56 (.51-.60)	.38 (.35-.42)	.41 (.37-.45)
No (reference)			
Dependent variable: Adequate physical activity			
Multimorbidity			
Yes	.56 (.53-.60)	.74 (.69-.80)	.82 (.76-.88)
No (reference)			
BRFSS 2017			
Dependent variable: Regular vegetable and/or fruit consumption			
Multimorbidity			
Yes	.91 (.88-.94)	.85 (.82-.88)	.91 (.88-.94)
No (reference)			

and policies promoting Whole Health activities must consider these nuanced relationships.

Our study results with respect to psychological distress are consistent with global studies in which multimorbidity has been found to be associated with psychological symptoms in 42 countries.<sup>37</sup> Although the causal pathways connecting multimorbidity to psychological stress are complex with numerous interacting factors such as lifestyle choices, and genetic, environmental, social, and behavioral factors,<sup>38</sup> our findings suggest monitoring mental health below the threshold for mental health disorders offers potential opportunities for early prevention.<sup>39</sup> In this context, the healthcare systems can provide support services to mitigate the effects of stress among adults with multimorbidity.

Our study found that those with multimorbidity were less likely to engage in recommended physical activities, even though associations with exercise and improvement of life expectancy in those with multimorbidity have been documented.<sup>40</sup> It is possible that pain and functional limitations may limit the participation of adults with multimorbidity in physical activities.<sup>41,42</sup> Low-impact activities such as water-based programs or stationary biking, or meditation-based activities such as yoga and Tai Chi are physical interventions that can be recommended for patients with chronic conditions if their abilities and conditions allow.<sup>43</sup>

Inadequate sleep has been linked to the development of chronic conditions such as diabetes, hypertension, and heart disease.<sup>44</sup> Moderate-severe sleep disturbances have also been associated with a higher speed of chronic disease accumulation, regardless of confounders.<sup>45</sup> A low percentage of adults with multimorbidity in our study had adequate sleep, similar to results found in prior studies. Additionally, dietary intake is central to overall health, and in our study, a lower proportion of adults with multimorbidity were less likely to report having regular vegetable and/or fruit consumption. Our findings are consistent with a cross-sectional study conducted by Dekker et al.,<sup>46</sup> where it was noted that higher adherence to a dietary pattern characterized by high intakes of meats, alcohol, potato, and also a “snack pattern” was associated with an increased prevalence of multimorbidity. Diet and sleep are closely related as studies have observed consistent associations with short sleep duration, higher caloric intake, higher absolute intake of fat, lower fruit intake, and lower quality diets. Therefore, Whole Health promotion strategies targeted at improving dietary intake and implementing cognitive behavioral therapy to help with insomnia should be considered in those with multimorbidity.

Previous research has documented a wide range of barriers to Whole Health activities among adults with multimorbidity.<sup>47</sup> These barriers suggest a multitiered and comprehensive approach such as the 1 proposed by the VA, where

a health system provides comprehensive support based on an individual's values, needs, and goals to promote Whole Health activities.<sup>7</sup> The comprehensive approach utilizes patient engagement support through education, wellness programs, and clinical care.<sup>7</sup>

Regarding study strengths, we used nationally representative data sets to analyze a comprehensive list of factors associated with Whole Health. This study also adds to the limited body of evidence on the direct analysis and prevalence of Whole Health activities. Those with multimorbidity had lower odds of adequate sleep, physical activity, regular consumption of vegetables and/or fruits, utilization of mind-body therapies, and low or no psychological distress. As published literature suggests that multimorbidity is associated with unfavorable clinical outcomes, integration and improvement of Whole Health activities for the prevention and management of multimorbidity may be beneficial.

Our study did have some notable limitations. This was a cross-sectional study design in which self-reported data is subject to recall data. A noteworthy finding is that we needed to analyze 2 different datasets to estimate the state of Whole Health self-care among adults in the U.S. Widespread adoption of the Whole Health model will require scientific evidence on its beneficial effects. Building such an evidence base is not possible without data on standardized measures of Whole Health. Unfortunately, beginning in 2019, NHIS has also moved away from collecting information on some of these measures every year. For example, data on psychological distress will be collected once in 3 years - 2019, 2022, and 2025. Thus, our findings suggest that research efforts and investment need to be made to collect nationally representative data on all Whole Health measures during the same time period. We support a call to action for collecting standardized Whole Health measures in datasets sponsored by federal agencies.

## Conclusion

Approximately one in 3 adults had multimorbidity. Adults with multimorbidity were less likely to report Whole Health activities, suggesting a system of support that moves away from a disease orientation to health promotion is needed. In addition, no 1 dataset collected information on all 5 activities of Whole Health activities during the same year. Therefore, we suggest collection of standardized measures on Whole Health activities for healthcare research, programs, and policies.

Based on 25 133 adult respondents (age 18 years or older) from the NHIS 2017 and 343 341 adult respondents from the BRFSS 2017. Model 1 adjusted only for multimorbidity variable. Model 2 adjusted for multimorbidity, sex, age, race and ethnicity, education, and poverty status. Model 3 is fully adjusted for multimorbidity, sex, age, race and ethnicity, education, poverty status, health insurance, marital status, body mass index, current smoking, alcohol use, and region.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: National Institute on Minority Health and Health Disparities, 5U54MD006882-10; National Institutes of Health, 1OT2OD032581-01 and National Institute of General Medical Sciences, 5U54GM104942-07.

## ORCID iD

Rolake A. Neba  <https://orcid.org/0000-0002-7348-6369>

## References

1. Koroukian SM, Warner DF, Owusu C, Given CW. Multimorbidity redefined: prospective health outcomes and the cumulative effect of co-occurring conditions. *Prev Chronic Dis.* 2015;12:E55. doi:10.5888/pcd12.140478
2. Boyd CM, Darer J, Boulton C, Fried LP, Boulton L, Wu AW. Clinical practice guidelines and quality of care for older patients with multiple comorbid diseases: implications for pay for performance. *JAMA.* 2005;294(6):716-724. doi:10.1001/jama.294.6.716
3. Mamdani M, Ching A, Golden B, Melo M, Menzefricke U. Challenges to evidence-based prescribing in clinical practice. *Ann Pharmacother.* 2008;42(5):704-707. doi:10.1345/aph.1K283
4. Whole person health: What You Need to Know. National Center for Complementary and Integrative Health. <https://www.nccih.nih.gov/health/whole-person-health-what-you-need-to-know>. Published 2021. Accessed June 11, 2022.
5. Pacific Institute for Research & Evaluation UoW-M. Passport to Whole Health. In: Transformation VOoPCCC, ed. 2020.
6. A Whole Person Approach to Health and Healing - NIWH Programs. <https://www.wholehealtheducation.com/a-whole-person-health-approach-to-health-and-healing/>. Published April 29, 2021. Accessed June 11, 2022.
7. Marchand WR, Beckstrom J, Nazarenko E, et al. The veterans health administration whole health model of care: early implementation and utilization at a large healthcare system. *Mil Med.* 2020;185(11-12):e2150-e2157. doi:10.1093/milmed/usaa198
8. Whole Health Institute. The Purpose. <https://www.wholehealth.org/>. Published 2021. Accessed June 11, 2022.
9. Whole Health. TakeCare. <https://takecare.org/>. Published June 29, 2021. Accessed June 11, 2022.
10. Integrative Medicine and the Health of the Public: A Summary of the February 2009 Summit. 2009.
11. Tripp-Reimer T, Williams JK, Gardner SE, et al. An integrated model of multimorbidity and symptom science. *Nurs Outlook.* 2020;68(4):430-439. doi:10.1016/j.outlook.2020.03.003

12. Crocker RL, Hurwitz JT, Grizzle AJ, et al. Real-world evidence from the integrative medicine primary care trial (IMPACT): assessing patient-reported outcomes at baseline and 12-month follow-up. *Evid Based Complement Alternat Med*. 2019;2019: 8595409. doi:10.1155/2019/8595409
13. Bokhour B, Hyde J, Zeliadt S, Mohr H. Whole health system of care evaluation—a progress report on outcomes of the WHS pilot at 18 Flagship sites. [(accessed on 11 June 2021)];Cent. Eval. Patient Cent. Care VA EPCC-VA. 2020 3.
14. Riegel B, Westland H, Iovino P, et al. Characteristics of self-care interventions for patients with a chronic condition: a scoping review. *Int J Nurs Stud*. 2021;116:103713. doi:10.1016/j.ijnurstu.2020.103713
15. Bokhour BG, Haun JN, Hyde J, Charns M, Kligler B. Transforming the veterans affairs to a whole health system of care: time for action and research. *Med Care*. 2020;58(4): 295-300. doi:10.1097/MLR.0000000000001316
16. Nelson EC, Meyer G, Bohmer R. Self-care: the new principal care. *J Ambul Care Manage*. 2014;37(3):219-225. doi:10.1097/JAC.0000000000000036
17. Sindi S, Perez LM, Vetrano DL, et al. Sleep disturbances and the speed of multimorbidity development in old age: results from a longitudinal population-based study. *BMC Med*. 2020; 18(1):382. doi:10.1186/s12916-020-01846-w
18. Bricca A, Harris LK, Jager M, Smith SM, Juhl CB, Skou ST. Benefits and harms of exercise therapy in people with multimorbidity: a systematic review and meta-analysis of randomised controlled trials. *Ageing Res Rev*. 2020;63:101166. doi: 10.1016/j.arr.2020.101166
19. Dankel SJ, Loenneke JP, Loprinzi PD. Combined associations of muscle-strengthening activities and accelerometer-assessed physical activity on multimorbidity: findings from NHANES. *Am J Health Promot*. 2017;31(4):274-277. doi:10.4278/ajhp.150520-QUAN-894
20. Dekker LH, de Borst MH, Meems LMG, de Boer RA, Bakker SJL, Navis GJ. The association of multimorbidity within cardio-metabolic disease components with dietary patterns: a cross-sectional study in 129 369 men and women from the Lifelines cohort. *PLoS One*. 2019;14(8):e0220368. doi:10.1371/journal.pone.0220368
21. Fortin M, Bravo G, Hudon C, Lapointe L, Dubois MF, Almirall J. Psychological distress and multimorbidity in primary care. *Ann Fam Med*. 2006;4(5):417-422. doi:10.1370/afm.528
22. Alwhaibi M, Bhattacharya R, Sambamoorthi U. Type of multimorbidity and complementary and alternative medicine use among adults. *Evid Based Complement Alternat Med*. 2015;2015:362582. doi:10.1155/2015/362582
23. Wahbeh H, Elsas SM, Oken BS. Mind-body interventions: applications in neurology. *Neurology*. 2008;70(24):2321-2328. doi:10.1212/01.wnl.0000314667.16386.5e
24. Watson NF, Badr MS, Belenky G, et al. Recommended amount of sleep for a healthy adult: a joint consensus statement of the American academy of sleep medicine and sleep research society. *Sleep*. 2015;38(6):843-844. doi: 10.5665/sleep.4716
25. Physical activity guidelines for Americans. *Okla Nurse*. 2009; 53(4):25.
26. Reed P. Physical activity: we all can and must do more. Physical Activity: We All Can and Must Do More - News and Events. <https://health.gov/news/202205/physical-activity-we-all-can-and-must-do-more-0>. Published May 18, 2022. Accessed June 11, 2022.
27. Andrews G, Slade T. Interpreting scores on the Kessler psychological distress scale (K10). *Aust N Z J Public Health*. 2001; 25(6):494-497. doi:10.1111/j.1467-842x.2001.tb00310.x
28. Prochaska JJ, Sung HY, Max W, Shi Y, Ong M. Validity study of the K6 scale as a measure of moderate mental distress based on mental health treatment need and utilization. *Int J Methods Psychiatr Res*. 2012;21(2):88-97. doi:10.1002/mpr.1349
29. Using the New BRFSS Module. Centers for Disease Control and Prevention. <https://www.cdc.gov/nutrition/data-statistics/using-the-new-BRFSS-modules.html>. Published September 1, 2020. Accessed June 11, 2022.
30. Salive ME. Multimorbidity in older adults. *Epidemiol Rev*. 2013;35:75-83. doi:10.1093/epirev/mxs009
31. Goodman RA, Posner SF, Huang ES, Parekh AK, Koh HK. Defining and measuring chronic conditions: imperatives for research, policy, program, and practice. *Prev Chronic Dis*. 2013;10:E66. doi:10.5888/pcd10.120239
32. Variance Estimation Guidance, NHIS 2016-2017. Centers for Disease Control and Prevention. <https://www.cdc.gov/nchs/data/nhis/2016var.pdf>. Published October 2018. Accessed June 11, 2022.
33. Iachan R, Pierannunzi C, Healey K, Greenlund KJ, Town M. National weighting of data from the behavioral risk factor surveillance system (BRFSS). *BMC Med Res Methodol*. 2016; 16(1):155. doi:10.1186/s12874-016-0255-7
34. Riegel B, Dunbar SB, Fitzsimons D, et al. Self-care research: where are we now? Where are we going? *Int J Nurs Stud*. 2021; 116:103402. doi:10.1016/j.ijnurstu.2019.103402
35. Malecki HL, Gollie JM, Scholten J. Physical activity, exercise, whole health, and integrative health coaching. *Phys Med Rehabil Clin N Am* 2020;31(4):649-663. doi:10.1016/j.pmr.2020.06.001
36. Black LI, Nugent CN, Adams PF. Tables of adult health behaviors, sleep: National Health Interview Survey, 2011–2014. In: Adult Health Behavior Tables: National Health Interview Survey -, Tables SLP-2CCdc-pdf, ed. 2016.
37. Vancampfort D, Koyanagi A, Hallgren M, Probst M, Stubbs B. The relationship between chronic physical conditions, multimorbidity and anxiety in the general population: a global perspective across 42 countries. *Gen Hosp Psychiatr*. 2017;45: 1-6. doi:10.1016/j.genhosppsy.2016.11.002
38. McEwen BS, Stellar E. Stress and the individual. Mechanisms leading to disease. *Arch Intern Med*. 1993;153(18): 2093-2101.
39. Arango C, Diaz-Caneja CM, McGorry PD, et al. Preventive strategies for mental health. *Lancet Psychiatr*. 2018;5(7): 591-604. doi:10.1016/S2215-0366(18)30057-9



40. Chudasama YV, Khunti KK, Zaccardi F, et al. Physical activity, multimorbidity, and life expectancy: a UK Biobank longitudinal study. *BMC Med.* 2019;17(1):108. doi:[10.1186/s12916-019-1339-0](https://doi.org/10.1186/s12916-019-1339-0)
41. Dorenkamp S, Mesters I, Vos R, et al. Synergistic effects of six chronic disease pairs on decreased physical activity: the SMILE cohort study. *BioMed Res Int.* 2016;2016:9427231. doi:[10.1155/2016/9427231](https://doi.org/10.1155/2016/9427231)
42. St John PD, Tyas SL, Menec V, Tate R, Griffith L. Multimorbidity predicts functional decline in community-dwelling older adults: prospective cohort study. *Can Fam Physician.* 2019;65(2):e56-e63.
43. Mora JC, Valencia WM. Exercise and older adults. *Clin Geriatr Med.* 2018;34(1):145-162. doi:[10.1016/j.cger.2017.08.007](https://doi.org/10.1016/j.cger.2017.08.007)
44. Dashti HS, Scheer FA, Jacques PF, Lamou-Fava S, Ordovas JM. Short sleep duration and dietary intake: epidemiologic evidence, mechanisms, and health implications. *Adv Nutr.* 2015;6(6):648-659. doi:[10.3945/an.115.008623](https://doi.org/10.3945/an.115.008623)
45. Sindi S, Perez LM, Vetrano DL, et al. Sleep disturbances and the speed of multimorbidity development in old age: results from a longitudinal population-based study. *BMC Med.* 2020;18(1):382. doi:[10.1186/s12916-020-01846-w](https://doi.org/10.1186/s12916-020-01846-w)
46. Dekker LH, de Borst MH, Meems LMG, de Boer RA, Bakker SJL, Navis GJ. The association of multimorbidity within cardio-metabolic disease components with dietary patterns: a cross-sectional study in 129 369 men and women from the Lifelines cohort. *PLoS One.* 2019;14(8):e0220368. doi:[10.1371/journal.pone.0220368](https://doi.org/10.1371/journal.pone.0220368)
47. Bayliss EA, Steiner JF, Fernald DH, Crane LA, Main DS. Descriptions of barriers to self-care by persons with comorbid chronic diseases. *Ann Fam Med.* 2003;1(1):15-21. doi:[10.1370/afm.4](https://doi.org/10.1370/afm.4)